#### EUROPEAN JOURNAL OF PURE AND APPLIED MATHEMATICS

2025, Vol. 18, Issue 2, Article Number 5924 ISSN 1307-5543 – ejpam.com Published by New York Business Global



# Computational Methods, Existence and Uniqueness for Solving 2-D Fractional Nonlinear Fredholm Integro-Differential Equation

Abeer M. Al-Bugami<sup>1,\*</sup>, Nuha A. Alharbi<sup>1</sup>, Amr M. S. Mahdy<sup>1</sup>

<sup>1</sup> Department of Mathematics and Statistics, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia

Abstract. In this paper, we have investigated the two-dimensional fractional nonlinear Fredholm integro-differential equation (**TDFNFI-DE**). These equations are used in many fields, including particle dynamics in physics, biology, and control theory. We have developed an effective combined approach in our work that uses Homotopy analysis (**HAM**) and the Adomian decomposition method (ADM) to solve fractional integro-differential equations in two dimensions. Numerical experiment results show the effectiveness of our recently created technique. We prove the existence and uniqueness of the exact solution. To illustrate the numerical effectiveness of the suggested approach is accurate and applicable to various nonlinear issues in science, according to theoretical and numerical results. **2020 Mathematics Subject Classifications**: 45J05, 45G10, 26A33, 35R11, 65R20, 65L10 **Key Words and Phrases**: TDFNFI-DE, existence and uniqueness, ADM, HAM

## 1. Introduction

Integral equations are used in many different fields, such as continuous mechanical engineering, potential concept, geophysics, electricity and magnetism, kinetic theory of gasses, hereditary phenomena in biology and physics, quantum mechanics, radiation, optimisation, which includes optimal system design, theory of communication, economics of mathematics, genetics of populations, medical procedures, computational problems of radiative equilibrium, the particle transport problems of the fields of a and reactor principle, acoustics, steady state heat conduction, and radiative heat transfer troubles. Fredholm integral equation is one of the most important integral equations. Singh and Pippal [1] utilized the Shehu transform combined with the Adomian polynomial to solve effectively nonlinear fractional differential equations. Additionally, Ali et al. [2] presented a numerical solution for fractional integro-differential equations with a linear functional argument,

*Email addresses:* abeer101aa@yahoo.com (A. Al-Bugami), noha1231@hotmail.com (N. Alharbi), amr\_mahdy85@yahoo.com (A. Mahdy)

1

https://www.ejpam.com

Copyright: (c) 2025 The Author(s). (CC BY-NC 4.0)

<sup>\*</sup>Corresponding author.

DOI: https://doi.org/10.29020/nybg.ejpam.v18i2.5924

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 2 of 14

employing Chebyshev serious for increased precision. Ovedepo et al. [3] contributed a competitional algorithm specifically designed for fractional Fredholm integro-differential equations, providing a robot and efficacious method for tackling these complex problems. Moreover, Toma and Postavaru [4] proposed a numerical method for solving fractional Fredholm integro-differential equations with a focus on enhancing accuracy. Abdou et al. is performed a comprehensive analysis of the convergence into that Adomian's method when applied to nonlinear equations in [5]. Abbaoui and Cherruault [6] also performed a comprehensive analysis of the convergence into that Adomian's method when applied to nonlinear equations. Behiry et al. [7] developed a new algorithm for the decomposition solution of nonlinear differential equations, further advancing the method. Tair et al. [8] explored two numerical techniques for solving the linear integro-differential Fredholm equation with a weakly singular kernel, while Al-Bugami [9] focused on two-dimensional Fredholm integro-differential equations with singular kernels, providing accurate numerical method for their solution. Abbaszadeh et al. [10] employed Legendre wavelets to solve fractional Fredholm integro-differential equations, while Rawashdeh et al. [11] provided a detailed convergence analysis for the fractional decomposition method as applied to nonlinear fractional Fredholm integro-differential equations. Li and Pang [12] expanded the application of the ADM to nonlinear systems. In addition, Momani and Noor [13] developed numerical techniques to solve fractional integro-differential equations of fourth order. In the realm of two-dimensional nonlinear Fredholm integro-differential equations, Al-Bugami [14] presented effective numerical solutions. Additionally, Hasan [15] applied the (ADM) to nonlinear systems involving fractional Fredholm integro-differential equations.

We will use the **HAM** and **ADM** for dealing with the **(TDFNFI-DE)** in this study. The TDFNFI-DE form will evaluate.

$$D^{\alpha}u(s,t) = f(s,t) + \lambda \int_{a}^{b} \int_{c}^{d} K(s,t,\tau,\xi)\gamma(\tau,\xi,D^{\alpha}u(\tau,\xi))d\tau d\xi,$$
(1)

where u(s,t) is an unknown function,  $f(s,t), K(s,t,\tau,\xi)$  are given functions, and  $\lambda$  is parameter. Also  $0 < \alpha \leq 1$  and  $(s,t) \in \Omega = [a,b] \times [c,d]$ .

## 2. The Existence and Uniqueness

**Theorem 1.** Let  $(\mathbf{M},d)$  be a metric space with  $\mathbf{M} \neq \emptyset$ . Suppose  $\mathbf{M} \neq \emptyset$  is complete and let  $I : \mathbf{M} \to \mathbf{M}$  be contraction mapping. Then, I has exactly one fixed point. Furthermore, we can express the equation in (1) in the form of an integral operator as follows:

$$\tilde{T}D^{\alpha}u(s,t) = f(s,t) + TD^{\alpha}u(s,t),$$
(2)

where

$$TD^{\alpha}u(s,t) = \lambda \int_{a}^{b} \int_{c}^{d} K(s,t,\tau,\xi)\gamma(\tau,\xi,D^{\alpha}u(\tau,\xi))d\tau d\xi.$$
(3)

Also assume the following conditions:

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 3 of 14

- (i)  $K(s, t, \tau, \xi)$  satisfies  $|K(s, t, \tau, \xi)| \leq \kappa$ , where  $\kappa$  is a constant.
- (ii) f(s,t) is continuous in  $C[a,b] \times C[c,d]$ , with its norm defined as:

$$||f(s,t)|| = \left[\int_{a}^{b} \int_{c}^{d} |f(s,t)|^{2} ds dt\right]^{\frac{1}{2}} = v,$$

where v is a constant.

(iii) The continuous function  $\gamma(s, t, D^{\alpha}u(s, t))$  satisfies the following conditions for some constant  $g > g_1 > m, g > m$ :

i. 
$$\left[\int_{a}^{b}\int_{c}^{d}|\gamma(s,t,D^{\alpha}u(s,t))|^{2}dsdt\right]^{\frac{1}{2}} \leq g_{1}||D^{\alpha}u(s,t)||.$$
  
ii. 
$$|\gamma(s,t,D^{\alpha}u_{1}(s,t))-\gamma(s,t,D^{\alpha}u_{2}(s,t))| \leq \mathbf{M}(s,t) \cdot |D^{\alpha}u_{1}(s,t)-D^{\alpha}u_{2}(s,t)|,$$
  
where  $\|\mathbf{M}(s,t)\| = M$  and  $M$  is a constant.

(iv) The unknown function  $D^{\alpha}u(s,t)$  behaves similarly to the given function f(s,t) in  $C[a,b] \times C[c,d]$ , with its norm defined as:

$$||D^{\alpha}u(s,t)|| = \left[\int_{a}^{b} \int_{c}^{d} |D^{\alpha}u(s,t)|^{2} ds dt\right]^{\frac{1}{2}}.$$

**Theorem 2.** If the first three conditions are satisfied, then eq (1) has a unique solution in  $C[a, b] \times C[c, d]$ .

**Lemma 1.** The space  $C[a,b] \times C[c,d]$  is mapped into itself by the operator  $\tilde{T}$  defined by (2) under the conditions (1)–(3-i).

*Proof.* Using equation (2) and (3), we have:

$$\tilde{T}D^{\alpha}u(s,t) \| \leq \|f(s,t)\| + |\lambda| \left\| \int_{a}^{b} \int_{c}^{d} |K(s,t,\tau,\xi)| |\gamma(\tau,\xi,D^{\alpha}u(\tau,\xi))| d\tau d\xi \right\|,$$
(4)

by applying condition (2), it follows that:

$$\|\tilde{T}D^{\alpha}u(s,t)\| \le v + |\lambda| \cdot \left[|K(s,t,\tau,\xi)|\right] \left[\int_{a}^{b} \int_{c}^{d} |\gamma(s,t,D^{\alpha}u(s,t))|^{2} ds dt\right]^{\frac{1}{2}},$$
(5)

using conditions (1) and (3-i), we can estimate:

$$\|\tilde{T}D^{\alpha}u(s,t)\| \le v + \delta \|D^{\alpha}u(s,t)\|,$$

when  $\delta = |\lambda| \kappa g$ . Hence, this inequality (2) confirms that the operator  $\tilde{T}$  is bounded, with:  $\|\tilde{T}D^{\alpha}u(s,t)\| \leq \delta \|D^{\alpha}u(s,t)\|.$  A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 4 of 14

**Lemma 2.** The operator  $\tilde{T}$  is a contraction on the Banach space  $C[a, b] \times C[c, d]$ , if conditions (1) and (3-ii) are satisfied.

*Proof.* Suppose we have two unknown function  $D^{\alpha}u_1(s,t)$  and  $D^{\alpha}u_2(s,t)$  in  $C[a,b] \times C[c,d]$ , we can apply equations (2) and (3) to obtain:

$$\begin{aligned} \|(\tilde{T}D^{\alpha}u_1 - \tilde{T}D^{\alpha}u_2)(s,t)\| &\leq |\lambda| \left\| \int_a^b \int_c^d |K(s,t,\tau,\xi)| \cdot |\gamma(\tau,\xi,D^{\alpha}u_1(\tau,\xi)) - \gamma(\tau,\xi,D^{\alpha}u_2(\tau,\xi))| d\tau d\xi \right\| \end{aligned}$$

by using condition (3-ii), we get:

$$\begin{aligned} \|(\tilde{T}D^{\alpha}u_1 - \tilde{T}D^{\alpha}u_2)(s,t)\| &\leq |\lambda| \cdot \left[|K(s,t,\tau,\xi)|\right] \cdot \\ \left[\int_a^b \int_c^d \mathbf{M}^2(s,t)|D^{\alpha}u_1(\tau,\xi) - D^{\alpha}u_1(\tau,\xi)|^2 d\tau d\xi\right]^{\frac{1}{2}}. \end{aligned}$$

Thus, we conclude:

$$\|(\tilde{T}D^{\alpha}u_{1} - \tilde{T}D^{\alpha}u_{2})(s,t)\| \le \delta \|D^{\alpha}u_{1}(s,t) - D^{\alpha}u_{2}(s,t)\|$$

#### **3.** ADM

In this section, we will address the **TDFNFI-DE** using ADM.

Let's consider (1), in witch f(s,t) is a function that is bounded, and for all (s,t) belongs to the set  $\Omega = [a,b] \times [c,d]$ . The kernel  $|K(s,t,\tau,\xi)| \leq \kappa$ . The nonlinear term  $\gamma(\tau,\xi, D^{\alpha}u(\tau,\xi))$  is Lipchitz continuous and satisfies the following condition:

$$|\gamma(D^{\alpha}u) - \gamma(D^{\alpha}\tilde{u})| \le L|D^{\alpha}u - D^{\alpha}\tilde{u}|.$$

We define the space  $C[a, b] \times C[c, d]$ , continuous functions on the rectangle  $[a, b] \times [c, d]$ , with the associated distance function  $d^*(D^{\alpha}\tilde{u}, D^{\alpha}u)$  defined by:

$$d^*(D^{\alpha}\tilde{u}, D^{\alpha}u) = \max_{(s,t)\in\Omega} |D^{\alpha}\tilde{u}(s,t) - D^{\alpha}u(s,t)|.$$
(6)

The unknown function  $D^{\alpha}u(s,t)$  is assumed to have the series form:

$$D^{\alpha}u(s,t) = \sum_{n=0}^{\infty} D^{\alpha}u_n(s,t).$$
(7)

The nonlinear term  $\gamma(\tau, \xi, D^{\alpha}u(\tau, \xi))$  in (1) is similarly decomposed into an infinite series:

$$\gamma(\tau,\xi,D^{\alpha}u(s,t)) = \sum_{n=0}^{\infty} A_n u_n(s,t),$$
(8)

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, **18** (2) (2025), 5924 5 of 14 where the traditional form of the Adomian polynomials  $A_n$  is [16][17] given by:

$$A_n = \frac{1}{n!} \left[ \frac{d^n}{d\lambda^n} \gamma \left( \sum_{i=0}^{\infty} \lambda^i D^\alpha u_i \right) \right]_{\lambda=0}.$$
 (9)

Another formulation of the Adomian polynomials is expressed as:

$$A_n = \gamma(S_n) - \sum_{i=0}^{n-1} A_i,$$
(10)

where  $S_n$  is defined by:

$$S_n = \sum_{i=0}^n D^{\alpha} u_i(s, t).$$
 (11)

By applying the Adomian decomposition method to equation (1), we obtain the solution in series form:

$$D^{\alpha}u(s,t) = \sum_{n=0}^{\infty} D^{\alpha}u_n(s,t), \qquad (12)$$

where

$$D^{\alpha}u_{0}(s,t) = f(s,t),$$
  

$$D^{\alpha}u_{i}(s,t) = \lambda \int_{a}^{b} \int_{c}^{d} K(s,t,\tau,\xi)A_{i-1}d\tau d\xi, \qquad i \ge 1.$$
(13)

# **4. HAM**

In this section, we aim to solve the nonlinear **(FNT-DFIDE)** using the Homotopy Analysis Method (HAM). This approach is novel because, to the best of our knowledge, previous studies have primarily applied HAM to solve two-dimensional fractional integrodifferential equations. Consider (1), where  $K(s, t, \tau, \xi)$  and f(s, t) are known functions, and  $\gamma(\tau, \xi, D^{\alpha}u(\tau, \xi))$  is a known nonlinear function of  $D^{\alpha}u$ . To describe the method, we define the nonlinear operator N:

$$N[D^{\alpha}u] = D^{\alpha}u(x,t) - f(x,t) - \int_{a}^{b} \int_{c}^{d} K(x,t,\tau,\xi)\gamma(\tau,\xi,D^{\alpha}u(\tau,\xi))d\tau d\xi = 0.$$
(14)

Let  $D^{\alpha}u_0(s,t)$  denote an initial guess for the exact solution u(s,t), and choose a non-zero auxiliary parameter  $h \neq 0$ , an auxiliary function H(s,t), and linear operator L, such that L[e(s,t)] = 0 whenever e(s,t) = 0. For  $p \in [0,1]$ , we construct the following homotopy:

$$(1-p)[\phi(s,t;p) - D^{\alpha}u_0(x,t)] - phH(s,t)N[\phi(s,t;p)] = H[\phi(s,t;p); D^{\alpha}u_0(s,t), H(s,t), h, p].$$
(15)

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 6 of 14

it is important to emphasize that we have significant flexibility in choosing the initial guess  $D^{\alpha}u_0(s,t)$ , the auxiliary linear operator L, the non-zero parameter h, and the auxiliary function H(s,t). Additionally, let  $\tilde{H}$  be a secondary auxiliary function that ensures the homotopy satisfies the condition:

$$H[\phi(s,t;p); D^{\alpha}u_0(s,t), H(s,t), h, p] = 0.$$
(16)

Thus, the homotopy equation can be written as:

$$(1-p)[\phi(s,t;p) - D^{\alpha}u_0(x,t)] = phH(s,t)N[\phi(s,t;p)],$$
(17)

when p = 0, from equation (17), we get:

$$\phi(s,t;p) = D^{\alpha}u_0(s,t). \tag{18}$$

Furthermore, when p = 1 and  $h \neq 0$ ,  $H(s, t) \neq 0$ , equation (17) becomes:

$$\phi(s,t;1) = D^{\alpha}u(s,t). \tag{19}$$

If we observe the value of p from 0 to 1m the embedding parameter gradually increases for  $\phi(s,t;p)$  in equations (1) and (19), such that it begins from the initial approximation and reaches the exact solution. We refer to this change in homotopy deformation, and we can be expressed in the following form:

$$\phi(s,t;p) = D^{\alpha}u_0(x,t) + \sum_{r=1}^{\infty} D^{\alpha}u_r(s,t)p^r.$$
 (20)

Here,  $u_r(s,t)$  is given by:

$$D^{\alpha}u_{r}(s,t) = \frac{1}{r!} \frac{\partial\phi(s,t;p)}{\partial p^{r}}\Big|_{p=0}$$
(21)

Based on these assumptions, we can write the exact solution as:

$$D^{\alpha}u(s,t) = \phi(s,t;1) = D^{\alpha}u_0(s,t) + \sum_{r=1}^{\infty} D^{\alpha}u_r(s,t)p$$
(22)

Thus, the sequence of approximations for  $D^{\alpha}u(s,t)$  can be expressed as:

$$D^{\alpha}\bar{u}_{n}(s,t) = \{D^{\alpha}u_{0}(s,t), D^{\alpha}u_{1}(s,t), \dots, D^{\alpha}u_{n}(s,t)\}$$
(23)

The zero-order deformation equation (21), in accordance with equation (17), can be used to construct the governing equation of  $D^{\alpha}u_m(s,t)$ . The called *m*th-order deformation equation gets by differentiating the zero-order deformation equation (17) r times with regard to p, dividing the result by r!, and setting r = 0.

Besed on equation (21), the governing equation for  $D^{\alpha}u_m(s,t)$  can be derived from the zero-order deformation equation (17). By differentiating equation (17) r times with A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, **18** (2) (2025), 5924 7 of 14 respect to r, then dividing by r! and setting p = 0, we obtain the *m*th-order deformation equation as follows:

$$L[D^{\alpha}u_{r}(s,t) - \sigma_{r}D^{\alpha}u_{r-1}(s,t)] = hH(s,t)\mathcal{J}(D^{\alpha}\bar{u}_{r-1}(s,t))$$
  
$$D^{\alpha}u_{r}(0,0) = 0$$
(24)

where

$$\mathcal{J}(D^{\alpha}\bar{u}_{r-1}(s,t)) = \frac{1}{(r-1)!} \frac{\partial^{r-1} N[\phi(s,t;p)]}{\partial p^{r-1}} \Big|_{p=0}$$
(25)

and the condition  $\sigma_r$  is defined as:

$$\sigma_r = \begin{cases} 0 & (r \le 0) \\ \\ 1 & (r > 0) \end{cases}$$
(26)

# 5. Numerical Applications

**Application 1.** Consider the following Caputo fractional linear Fredholm integro-differential equation

$$D^{0.5}u(s,t) = 2.256758334 \ s\sqrt{t} - 0.6790610904 + \int_0^1 \int_0^1 \sqrt{\tau\xi} \ D^{0.5}u(\tau,\xi)d\tau d\xi \tag{27}$$

The exact solution  $u(s,t)=2.256758334\;s\sqrt{t}$  .

Table 1: Numerical results and absolute error values using (ADM) and (HAM), N = 10, at fractional order  $\alpha = 0.5$  in Application 1.

s	t	$u_{Exact}$	ADM		НАМ		
			$u_{ADM}$	$Error_{ADM}$	$u_{HAM}$	$Error_{HAM}$	
0.0	0.0	0	0	0	$4 \times 10^{-8}$	$4 \times 10^{-8}$	
0.1	0.1	0.07136496464	0.07136496464	0	0.07136500464	$4.000 \times 10^{-8}$	
0.2	0.2	0.2018506018	0.2018506017	$1 \times 10^{-10}$	0.2018506417	$3.99  imes 10^{-8}$	
0.3	0.3	0.3708232338	0.3708232338	0	0.3708232738	$4.00 \times 10^{-8}$	
0.4	0.4	0.5709197172	0.5709197171	$1 \times 10^{-10}$	0.5709197571	$3.99  imes 10^{-8}$	
0.5	0.5	0.797884561	0.7978845608	$2 \times 10^{-10}$	0.7978846008	$3.98 \times 10^{-8}$	
0.6	0.6	1.048846493	1.048846493	0	1.048846533	$4.0 \times 10^{-8}$	
0.7	0.7	1.321697642	1.321697641	$1 \times 10^{-9}$	1.321697681	$3.9 \times 10^{-8}$	
0.8	0.8	1.614804814	1.614804814	0	1.614804854	$4.0 \times 10^{-8}$	
0.9	0.9	1.926854045	1.926854045	0	1.926854085	$4.0  imes 10^{-8}$	
1.0	1.0	2.256758334	2.256758334	0	2.256758374	$4.0  imes 10^{-8}$	

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 8 of 14



Figure 1: Exact Solution of (ADM) for  $\alpha = 0.5$  in Application 1



Figure 3: Exact Solution of (HAM) for  $\alpha = 0.5 \text{ in Application 1}$ 



Figure 2: Approximate Solution of (ADM) for  $\alpha = 0.5$  in Application 1



Figure 4: Approximate Solution of (HAM) for  $\alpha = 0.5$  in Application 1

**Application 2.** Consider the following Caputo fractional linear Fredholm integro-differential equation

$$D^{0.8}u(s,t) = 2.178248842 \ st^{1/5} - 0.8324154416 + \int_0^1 \int_0^1 \sqrt{\tau\xi} \ D^{0.8}u(\tau,\xi)d\tau d\xi \qquad (28)$$

The exact solution is  $u(s,t) = 2.178248842 \ st^{1/5}$ .

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 9 of 14

s	t	$u_{Exact}$	ADM		HAM	
			$u_{ADM}$	$Error_{ADM}$	$u_{HAM}$	$Error_{HAM}$
0.0	0.0	0	0	0	$3 \times 10^{-8}$	$3 \times 10^{-8}$
0.1	0.1	0.1374382105	0.1374382105	0	0.1374382405	$3.00  imes 10^{-8}$
0.2	0.2	0.3157500926	0.3157500925	$1 \times 10^{-10}$	0.3157501225	$2.99\times 10^{-8}$
0.3	0.3	0.5136330933	0.5136330933	0	0.5136331233	$3.00  imes 10^{-8}$
0.4	0.4	0.725403224	0.7254032241	$1 \times 10^{-10}$	0.7254032541	$3.01 \times 10^{-8}$
0.5	0.5	0.948137878	0.9481378781	$1 \times 10^{-10}$	0.9481379081	$3.01 \times 10^{-8}$
0.6	0.6	1.180018979	1.180018979	0	1.180019009	$3.0 \times 10^{-8}$
0.7	0.7	1.419793357	1.419793357	0	1.419793387	$3.0 \times 10^{-8}$
0.8	0.8	1.666538980	1.666538980	0	1.666539010	$3.0  imes 10^{-8}$
0.9	0.9	1.919545908	1.919545908	0	1.919545938	$3.0  imes 10^{-8}$
1.0	1.0	2.178248842	2.178248842	0	2.178248872	$3.0  imes 10^{-8}$

Table 2: Numerical results and absolute error values using (ADM) and (HAM), N = 10, at fractional order  $\alpha = 0.8$  in Application 2.



Figure 5: Exact Solution of (ADM) for  $\alpha = 0.8 \text{ in Application } 2$ 



Figure 6: Approximate Solution of (ADM) for  $\alpha = 0.8$  in Application 2

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 10 of 14







Figure 8: Approximate Solution of (HAM) for  $\alpha = 0.8$  in Application 2

**Application 3.** Consider the following Caputo fractional nonlinear Fredholm integrodifferential equation

$$D^{0.5}u(s,t) = 2.256758334 \ s\sqrt{t} - 1.235153476 + \int_0^1 \int_0^1 \sqrt{\tau\xi} \ (D^{0.5}u(\tau,\xi))^2 \ d\tau d\xi \quad (29)$$

s	t	$u_{Exact}$	ADM		$H\overline{A}M$	
			$u_{ADM}$	$Error_{ADM}$	$u_{HAM}$	$Error_{HAM}$
0.0	0.0	0	0	0	$-2 \times 10^{-8}$	$2 \times 10^{-8}$
0.1	0.1	0.07136496464	0.07136496464	0	0.07136494464	$2.000\times 10^{-8}$
0.2	0.2	0.2018506018	0.2018506017	$1 \times 10^{-10}$	0.2018505817	$2.01 \times 10^{-8}$
0.3	0.3	0.3708232338	0.3708232338	0	0.3708232138	$2.00 \times 10^{-8}$
0.4	0.4	0.5709197172	0.5709197171	$1 \times 10^{-10}$	0.5709196971	$2.01 \times 10^{-8}$
0.5	0.5	0.797884561	0.7978845608	$2 \times 10^{-10}$	0.7978845408	$2.02\times10^{-8}$
0.6	0.6	1.048846493	1.048846493	0	1.048846473	$2.0 \times 10^{-8}$
0.7	0.7	1.321697642	1.321697641	$1 \times 10^{-9}$	1.321697621	$2.1  imes 10^{-8}$
0.8	0.8	1.614804814	1.614804814	0	1.614804794	$2.0  imes 10^{-8}$
0.9	0.9	1.926854045	1.926854045	0	1.926854025	$2.0  imes 10^{-8}$
1.0	1.0	2.256758334	2.256758334	0	2.256758314	$2.0 \times 10^{-8}$

The exact solution is  $u(s,t) = 2.256758334 \ s\sqrt{t}$ .

Table 3: Numerical results and absolute error values using (ADM) and (HAM), N = 10, at fractional order  $\alpha = 0.5$  in Application 3.

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 11 of 14



Figure 9: Exact Solution of (ADM) for  $\alpha=0.5$  in Application 3



0.4

Figure 11: Exact Solution of (HAM) for  $\alpha = 0.5$  in Application 3

1.5

1-0.5

0.4



Figure 10: Approximate Solution of (ADM) for  $\alpha = 0.5$  in Application 3



Figure 12: Approximate Solution of (HAM) for  $\alpha = 0.5$  in Application 3

Application 4. Consider the following Caputo fractional nonlinear Fredholm integrodifferential equation

$$D^{0.8}u(s,t) = 2.178248842 \ st^{1/5} - 1.631364025 + \int_0^1 \int_0^1 \sqrt{\tau\xi} \ (D^{0.8}u(\tau,\xi))^2 \ d\tau d\xi \quad (30)$$

The exact solution is  $u(s,t) = 2.178248842 \ s \ t^{1/5}$ .

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 12 of 14

6	t	$u_{Exact}$	ADM		HAM	
			$u_{ADM}$	$Error_{ADM}$	$u_{HAM}$	$Error_{HAM}$
0.0	0.0	0	0	0	$-5.5 \times 10^{-8}$	$5.5  imes 10^{-8}$
0.1	0.1	0.1374382105	0.1374382105	0	0.1374381555	$5.50  imes 10^{-8}$
0.2	0.2	0.3157500926	0.3157500925	$1 \times 10^{-10}$	0.3157500375	$5.51  imes 10^{-8}$
0.3	0.3	0.5136330933	0.5136330933	0	0.5136330383	$5.50 imes10^{-8}$
0.4	0.4	0.725403224	0.7254032241	$1 \times 10^{-10}$	0.7254031691	$5.49  imes 10^{-8}$
0.5	0.5	0.948137878	0.9481378781	$1 \times 10^{-10}$	0.9481378231	$5.49  imes 10^{-8}$
0.6	0.6	1.180018979	1.180018979	0	1.180018924	$5.5  imes 10^{-8}$
0.7	0.7	1.419793357	1.419793357	0	1.419793302	$5.5  imes 10^{-8}$
0.8	0.8	1.66653898	1.66653898	0	1.666538925	$5.5  imes 10^{-8}$
0.9	0.9	1.919545908	1.919545908	0	1.919545853	$5.5  imes 10^{-8}$
1.0	1.0	2.178248842	2.178248842	0	2.178248787	$5.5 imes10^{-8}$

Table 4: Numerical results and absolute error values using (ADM) and (HAM), N = 10, at fractional order  $\alpha = 0.8$  in Application 4.



Figure 13: Exact Solution of (ADM) for  $\alpha = 0.8$  in Application 4



Figure 14: Approximate Solution of (ADM) for  $\alpha = 0.8$  in Application 4

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 13 of 14



Figure 15: Exact Solution of (HAM) for  $\alpha = 0.8$  in Application 4



Figure 16: Approximate Solution of (HAM) for  $\alpha = 0.8$  in Application 4

### 6. Conclusion

From the previous study, we concluded the following: as s, t are increasing in  $[0, 1] \times [0, 1]$  the error for (ADM) is increasing and for (HAM) is remain constant at 108. In nonlinear case the using 0.8 is a smaller than by 0.5. The error using 0.8 is it smaller than in (ADM) and (HAM),(ADM) using 0.8 is more accurate than using 0.5. At N = 10 the error is decreasing in linear and nonlinear case.When using the same fractional order (ADM) and (HAM), exactly the same error, but using fractional order 0.8, (ADM) convergences faster than (HAM). The error decreases when the fractional order of (ADM) increases and the error for (HAM) remains the same. When the fractional order of (ADM) is 0.8, the error increases, whereas when the fractional order is 0.8, the error decreases. The Maple program was used to write the codes. Future Work. Other methods, such as the Homomtopy perturbation method, the Galerkin method, the collocation method and Chypechev polynomial, will be used to solve the **TDFNFI-DE**, and **compare** with ADM and HAM.

#### References

- [1] Amandeep Singh and Sarita Pippal. Solving nonlinear fractional differential equations by using shehu transform and adomian polynomials. pages 797–816, 2024.
- [2] Khalid K Ali, Mohamed A Abd El Salam, Emad MH Mohamed, Bessem Samet, Sunil Kumar, and MS Osman. Numerical solution for generalized nonlinear fractional integro-differential equations with linear functional arguments using chebyshev series. 2020:494, 2020.
- [3] Taiye Oyedepo, Christie Yemisi Ishola, Adam Ajimoti Ishaq, Abdullahi Muhammed

A. M. Al-Bugami, N. A. Alharbi, A. M. S. Mahdy / Eur. J. Pure Appl. Math, 18 (2) (2025), 5924 14 of 14

Ayinde, and Olalekan Lukman Ahmed. Computational algorithm for fractional fredholm integro-differential equations. 17(1), 2023.

- [4] Antonela Toma and Octavian Postavaru. A numerical method to solve fractional fredholm-volterra integro-differential equations. 68:469–478, 2023.
- [5] M A Abdou, I L El-Kalla, and A M Al-Bugami. New approach for convergence of the series solution to a class of nonlinear hammerstein integral equations. 3(4):261–269, 2011.
- [6] K. Abbaoui and Y. Cherruault. Convergence of adomian's method applied to nonlinear equations. 20(9):69–73, 1994.
- [7] S.H. Behiry, H. Hashish, I.L. El-Kalla, and A. Elsaid. A new algorithm for the decomposition solution of nonlinear differential equations. 54(4):459–466, 2007.
- [8] Boutheina Tair, Sami Segni, Hamza Guebbai, and Mourad Ghait. Two numerical treatments for solving the linear integro-differential fredholm equation with a weakly singular kernel. 23(2):117–136, 2022.
- [9] Abeer M. Al-Bugami. Two-dimensional fredholm integro-differential equation with singular kernel and its numerical solutions. 2022(2501947):1–8, 2022.
- [10] D Abbaszadeh, M Tavassoli Kajani, M Momeni, M Zahraei, and M Maleki. Solving fractional fredholm integro-differential equations using legendre wavelets. 166:168– 185, 2021.
- [11] Mahmoud S Rawashdeh, Hala Abedalqader, and Nazek A Obeidat. Convergence analysis for the fractional decomposition method applied to class of nonlinear fractional fredholm integro-differential equation. 17:174830262211511, 2023.
- [12] Wenjin Li and Yanni Pang. Application of adomian decomposition method to nonlinear systems. 2020(1):67, 2020.
- [13] Shaher Momani and Muhammad Aslam Noor. Numerical methods for fourth-order fractional integro-differential equations. 182(1):754–760, 2006.
- [14] A. M. Al-Bugami. Nonlinear fredholm integro-differential equation in two-dimensional and its numerical solutions. 6(10):10383–10394, 2021.
- [15] Nabaa N Hasan. Adomian decomposition method applied to nonlinear system of fractional fredholm integro-differential equations. 24(5), 2013.
- [16] I.L. El-Kalla. New results on the analytic summation of adomian series for some classes of differential and integral equations. 217(8):3756–3763, 2010.
- [17] I.L. El-Kalla. Piece-wise continuous solution to a class of nonlinear boundary value problems. 4(2):325–331, 2013.